

## ON THE OLDEST ARCHIDISKODON ELEPHANT FROM THE ANTHROPOGENE OF THE USSR

V. E. GARUTT,

L. I. ALEXEJEVA,

V. S. BAIGUSHEVA,

ZOOLOGICAL INSTITUTE  
AC. SCI. USSR

GEOLOGICAL INSTITUTE  
AC. SCI. USSR

ROSTOV STATE  
UNIVERSITY.

### ABSTRACT

*A. gromovi* is one of the most primitive species among the elephants of the genus *Archidiskodon* which lived on the territory of Eurasia in the Villafranchian time. In the teeth structure of this elephant constant premolar  $p^4$  is preserved. The last true molar  $M^3$  consists, on the average, of 13 laminae (optimal number). Remains of *A. gromovi* occur in the southern part of the USSR (from Moldavia to West Siberia and South of the Middle Asia). Geological age of the species: Upper Akchagyl (=Middle Villafranchian of the Western Europe).

Almost half a century ago remains of the ancient elephant were found in the sand-pit near Khapry station (suburb of the Rostov-on Don city) and preliminary referred to the *Elephas meridionalis* (Bogachev, 1924). Later on the Khapry sand formation and mammals fauna confined to it was studied by V. I. Gromov (1933, 1948). This scientist spoke out the assumption that the remains belong to two species, to the very archaic *Elephas planifrons* and to more progressive *E. meridionalis*. The skull of a young elephant found in Khapry sand-pit in 1934 is extremely interesting (Fig. 1, 2). Presence in this skull of a constant premolar  $P^4$  reasoned V. I. Gromov's assumption that the archaic elephant from Khapry sands is close to or identical with the *Protelephas planifrons* known from the Siwalik deposits of India.

The study of the following finds from Khapry and adjacent Liventsovka sand-pits proved the remains from Khapry sands to belong to one and the same species (Yan'kova, 1959). This elephant cannot be identical with the Indian *Protelephas planifrons* (Garutt, 1957). The elephant from Khapry sand-pit is in the skull structure a typical archidiskodont (its forehead is lengthwise strongly concave). Structural peculiarities of the skull and teeth allowed to consider the Khapry elephant an independent species which was named after the well-known Soviet geologist and paleontologist V. I. Gromov.

### SYSTEMATIC DESCRIPTION

Genus *Archidiskodon* Pohlig, 1885

*Archidiskodon gromovi* Garutt et Alexejeva, 1964

*Elephas meridionalis* Bogachev, 1923-1924, p. 108, Fig. 3-5; Gromov, 1933, p. 46; Gromov and Mirchink, 1936, p. 639, Fig. 174-K; Gromov, 1948, p. 44, 451-453, Fig. 214-C.

*Elephas planifrons* Gromov, 1933, p. 46; Gromov and Mirchink, 1936, p. 639, Fig. 174-I; Gromov, 1948, p. 44, 451-453, Fig. 214-B.

*Archidiskodon planifrons* Nikiforova and Alexejeva, 1959, p. 15, Table I.

*Archidiskodon meridionalis* Nikiforova and Alexejeva, 1959, p. 15, Table I; Dubovo and Baigusheva, 1964, pp. 133-136; Dubrovo 1964, pp. 86-88.

*Archidiskodon aff. meridionalis* Baigusheva, 1964, pp. 44-50.

*Archidiskodon gromovi* Garutt and Alexejeva, 1964, p. 78; Alexejeva and Garutt, 1965, pp. 171-175, Fig. I; Alexejeva, 1965, pp. 73-75, Table III-IV; 1970, pp. 124-125, Fig. 3; Baigusheva, 1968, pp. 5-29, b, Table III, Fig. 3-5; Vislobokova, 1974, pp. 173-175.

**Holotype:** (Pl. I). The skull of an old male (?) (without lower jaw and tusks) with last strongly worn out molar  $M^3$  from Khapry sands of the Liventsovka pit (Rostov-on Don city). Late Pliocene (early Eopleistocene according to V. I. Gromov's scheme of 1957). Museum of Regional Study in Rostov-on Don, N 113.

**Diagnosis:** The skull relatively low, longitudinally (sagittally) elongated: the ratio of its length (distance from the most protruding point on the face in its orbital area to Condylus occipitales inclusively) to the height (distance from molars' masticatory surface to the skull top) makes up 86 per cent. Occipital tubers (paired symmetrical protuberances on the occipital surface located slightly above Foramen occipitale magnum) are feebly prominent. Occipital surface lies at a right angle to the length axis of the skull (line running across the middle of the orbit and lower margin of Foramen acusticus externus). Forehead is narrow: the ratio of minimal forehead width to the skull width in its occipital region makes up 29 per cent. The lower jaw, mandibula, is rather low, for-

wards—backwards elongated. There is a strongly developed restrum. Teeth ontogenesis is characterized by the presence of a constant premolar P<sup>4</sup>. The last molar, M<sup>8</sup>, has the following (more frequently observed) characteristics: number of crown laminas—13; average length of one lamina—20 mm (on the upper teeth) and 22 mm (on the lower teeth); Lamina frequency in 10 cm of the crown—5.0 (on the upper teeth) and 4,5—4,8 (on the lower teeth); enamel thickness—2,9—3,9 mm.

*Description and comparison:* The skull of Gromov's elephant, when compared with that of *A. meridionalis*, shows well marked tendency to gradual increasing the height and simultaneous reducing its length (Fig. 1). In the holotype *A. gromovi* this ratio makes up 86 per cent, and in *A. meridionalis* it is slightly different; in the skull from Georgievka pit in the Northern Caucasus (museum of Regional Study in Pyatigorsk) the pointed

index constitutes 79,9 per cent; in the specimen from Alexandria pit, Northern Caucasus, (Zoological Museum, Ac. Sci. USSR, N 31065) it makes up 71,2 per cent and in the elephant skull from Taribana, Georgia, (Institute of Palaeobiology, Ac. Sci. of Georgian SSR, N k/31) —78, 0 per cent.

Occipital tubers are in *A. gromovi* of negligible prominence, the occipital region lying, therefore, at an almost right angle to the length axis of the skull. Occipital plane in *A. meridionalis* forms with the length axis of the skull an obtuse angle. In virtue of the latter the skull dome is strongly bent backwards (Weithofer, 1890, Table II, Fig. I); Maccagno, 1962, Fig. 5, Table II, Fig. 2; Gabunia and Vekua, 1963, Table II, Fig. 7. This peculiar skull structure in southern elephants seems to be connected with their more strongly developed occipital tubers.




Fig. 1. Skull of a young elephant of *Archidiskodon gromovi* from Khapry, side view. Coll. GIN, N 300—122.

TABLE I  
Skull measurements in *Archidiskodon gromovi*

NN	Measuring (in mm)	Liventsovka sandpit (c. Rostov-on-Don), Rostov Museum of Regional Study, N 112
1.	Distance from the skull top to anterior ends of intermaxillare .. ..	Ca 1370
2.	Distance from condyli occipitales to anterior ends of intermaxillare .. ..	1145
3.	Length of skull (distance from the front of the orbit to condyli occipitales) .. ..	805
4.	Height of skull (distance from molars' masticatory surface to the skull top) .. ..	Ca 934
5.	Occipital width of skull .. ..	895
6.	Width of skull at processus zygomaticus squamosum .. ..	835
7.	Height of occiput (from the skull top to condyli occipitales inclusively) .. ..	704
8.	Distance between the outer margins of condyli occipitales .. ..	228
9.	Size of foramen occipitales magnum (height × width) .. ..	81 × 80
10.	Length of forehead (from the skull top to the upper margin of intermaxillare) .. ..	540
11.	Minimal width of forehead .. ..	260
12.	Width of nasal hole .. ..	500
13.	Maximum height of nasal hole .. ..	135
14.	Length of intermaxillare .. ..	845
15.	Width of intermaxillare at the level of foramen infraorbitale .. ..	486
16.	Width of intermaxillare in the mid part .. ..	410
17.	Width of intermaxillare at alveolar margins of tusks .. ..	490
18.	Diameter of tusks .. ..	218 × 163
19.	Length of zygomatic arch .. ..	501
20.	Distance between the molars:	
	(a) from the front .. ..	135
	(b) from behind .. ..	204

In Gromov's elephant the forehead is sharply narrowed in its middle part, this narrowing being much more prominent than in *A. meridionalis*. Thus, in the holotype *A. gromovi* from Liventsovka sand-pit the ratio of the minimal forehead width to the skull width in its occipital portion makes up 29.0 per cent; in *A. meridionalis* from

Seneze (Basel Museum) —45.4 per cent, in that from Georgievka pit—45.3, from Alexandria—40.3 and from Taribans—47.5 per cent.

*A. gromovi* differs from the other species of the *Archidiskodon* genus also in the structure of its teeth row. The most essential feature proving the primitive level of Gromov's elephant is the presence of a constant premolar in its ontogenesis (Fig. 2). The study of mandibles in *A. meridionalis* specimens has shown that the constant premolars were absent in these elephants (Friant, 1959; Schaub, 1948 and others). The complete teeth series of *A. gromovi* could be designated as follows: 12, dP2, dP3, dP4, P4, M1, M2, M3.

Constant premolar tooth found in the skull of a young specimen of *A. gromovi* from Khapry (Gromov, 1948, Věrescagin, 1959, Garutt and Alexejeva 1964, Alexejeva and Garutt 1965, etc.)<sup>1</sup> is the last premolar P4. It sits directly before the functioning M<sup>1</sup> whose attachment to this succession is confirmed by the forming M<sup>2</sup> present in the mandibula.

P4 (Fig. 2) is similar in its crown structure to the constant premolars of mastodons. The crown is formed of some mammiform tubercles grouped in two indistinct rows. Both ends of the crown the tubercles are smaller and form the anterior and posterior talons. The enamel is vertically folded. Masticatory surface is covered with a dense cement layer. The tooth is not worn out. It seems not to be involved in masticatory process (first molar sitting behind P<sup>4</sup> is distinctly worn out). The described tooth (collection GIN, Ac. Sci. USSR, N 300-122) has the following size characteristics: length of crown—28 mm, width—32 mm, height—22 mm.

Molars of *A. gromovi* (in particular fragmentary molars) are very similar to those of *A. meridionalis*. To a certain degree they differ only in the last molars. M3 consists of 12-(13)-14 complete laminae. It should be noted that in the last molars found in Khapry and Liventsovka pits the number of complete laminae equals to 12-13 (not more). In *A. meridionalis* the number of laminae forming the crown is usually larger. It even amounts 16-17 laminae (Dubrovo, 1964).

The teeth of *A. gromovi* and those of *A. meridionalis* are almost of the same size. In *A. gromovi* the length of M3 crown is limited to the range of 200-320 mm; its width—to 80-120 mm and height—to 105-150 mm. In *A. meridionalis*, however, average height indices of the crown of M3 are slightly higher (sometimes they approach to 180-190 mm). In *A. gromovi* maximum width of the crown is at its base. Laminae are far removed from each other (there are 3,5—5,0) laminae in every

<sup>1</sup>Detailed description of this unique find presented by Prof. V. I. Gromov is now in press.




Fig. 2. Skull of a young elephant of *Archidiskodon gromovi* from Shapry, view from below.

100 mm of the crown length). In some cases there is a mid expansion on the upper teeth (especially on strongly worn out teeth). The pattern of laminae's abrasion is of mixed type. In the early stages of abrasion the antiquoid type was prevailing (side figures are rounded and the middle figure, elongated). In the middle stage of abrasion the meridianoid type is more frequently observed (side figures are oval and the middle figure, rounded). Sometimes figures both of the first and second types of abrasion are observed on one and the same tooth. The partition of tooth laminae into separate conelets proceeds rather deep (almost as deep as  $1/3$  of the lamina height). It is worth noting that the cement layer on the teeth of *A. gromovi* is developed to a lesser degree than on those of *A. meridionalis*. In certain places it does not completely fill the intervals between laminae (Plate II—1, 2).

*A. gromovi* is well distinguished from more late representatives of the *Archidiskodon* genus known from Europe by a number of structural peculiarities of the last molar. Thus, the crown of M3 in the elephants of the *trogontherii* species bears 18-20 complete laminae. In addition, these late elephant specimens always have teeth with more thin enamel (2-3 mm) and more closely set laminae (usually 5,5—6,5 in 100 mm of the crown length).

Gromov's elephant differs from *A. africanus* described from the Lower Villa-franchian of North Africa (Arambourg, 1952, 1970) in a large number of laminae on M3. In *A. africanus* the crown of a last molar bears 10 complete laminae (3,5 laminae in 100 mm).

From the North-American elephant specimens, which H. F. Osborn (1942) referred to the *Archidiskodon* genus, *A. gromovi* is well distinguished by the number of laminae

on M3 (in the North-American forms it is not known to be less than 17).

*Occurrence, age and habitat:* *Archidiskodon gromovi* was first identified by the remains dug out from Khapry sands (Gromov, 1933, 1948; Khokhlovkina, 1940). Their stratotype section is in Khapry sand-pit, near the city Rostov-on Don. At present this pit is not working. The Khapry formation is made up of alluvial deposits of one sedimentation cycle (Rodzianko, 1970). They occur along the northern coast of the Sea of Azov and at the Lower Don. This area was a field of acting of a river stream which ran along the Don valley. At present the Khapry sand formation is well exposed in Liventsovka pit (suburb of the city Rostov-on Don) whose section is considered to be the parastratotype of the Khapry formation. Khapry formation is 18 m thick. The floor of this old terrace is composed of marine deposits of Sarmatian, Meotian and Pontian age. Khapry formation is overlain by the thick series of Scyphian clays (Rodzianko, 1967). Bones accumulations of large mammals are confined to the lower 1/3 of this formation. Besides numerous elephant remains, the bones of *Equus livenzovensis* (= *E. cf. bressanus*), *E. ex gr. stenonis*, *Dicerorhinus etruscus*, *Elasmotherium caucasicum*, *Paracamelus alutensis* (= *P. kujalnensis*) *Eucladoceros dicranios*, *Gazellospira torticornis*, etc. have been found there too. Ostrich bones often occur in these bone-bearing lenses.

Fauna obtained from the Khapry and Liventsovka pits belongs to the Khaprovian faunistic complex and represents its early evolutionary stage (Alexejeva, 1968; Alexejeva, 1970). This part of the complex is correlating with the Middle Villafranchian of Europe (Alexejeva, 1974).

Fairly many localities of *A. gromovi* are known on the European territory of the USSR: in Moldavia—loc. Farladany, in Ukraine—Kotlovina; in the Rostov area—Khapry, Liventsovka, Marine, Valovaya Balka, Gunyushki, Chernorechye, Sablinskoe, etc. The tooth found in the Kosiakian pit (Gabunia, 1961) containing fauna of the Moldavian complex seems also probable to belong to that same elephant species. It is not excluded that the part of remains found on the territory of Azerbaijan SSR and identified as *Elephas planifrons* (Burchak-Abramovich and Djafarov, 1950) belongs to *A. gromovi*. It appears quite possible on account of the remains of *A. cf. gromovi* unearthed in the middle—Akchagyl part of the locality Kushkuna, Azerbaijan SSR (Lebedeva 1972). On the East the areal distribution of *A. gromovi* extends to the South of West Siberia (Vislobokova, 1974) and on the South its boundary approaches to South Tajikistan (Nikonov, Penkov *et al.*, 1971). The western boundary of its expansion is still not finally traced. In Western Europe teeth of primitive elephants were found

at some localities in the Lower Villafranchian deposits (Azzaroli, 1970). Their definite specific affinity so, as that of a tooth from the Kosiakian pit, is not determined yet. It seems not excluded that these remains belong to another even more ancient elephant specimen of the *Archidiskodon* genus.

*A. gromovi*, apparently, lived under conditions of the forest-shaped landscape, similar to savanna, and was mainly feeded on foliage, branches and tree bark. Eminent height of this elephant is due to adaptation for picking up food items from the large trees. Judging on the associating fauna composition, the climate was at that time rather warm, with hot summer and not heavy low-snow or snowless winter periods.


#### CONCLUSION



Recent data available on the occurrence of the remains of ancient *Archidiskodon* elephants permit the assertion that in the time of existence of fauna of early Khaprovian complex (upper part of the Akchagyl stage) *A. gromovi* was widely spread in warm latitudes of the North Eurasia. Its teeth structure is characterized by the presence of a last constant premolar. It makes us to assume that the above species could be a contemporary of the Indian *Protelephas planifrons*, in whose ontogenesis presence of the last constant premolars is also known. *A. gromovi* appears to be the direct ancestor of *A. meridionalis* (Garutt, 1964). Elephantine teeth found in the deposits somewhat older than that of the Khapry locality indicate that the *Archidiskodon* elephants appeared in Europe as a component of the Moldavian faunal complex which is correlating with the Rousilion or Villafranchian of the Western Europe (Alexejeva, 1974).

According to the wish of Acad. Yu. A. Orlov the skull of a young elephant of *A. gromovi* from Khapry (as well as that of an adult elephant from Liventsovka) will be given over to the Palaeontological Institute, Ac. Sci. USSR, for exhibition in the new Palaeontological Museum named after Yu. A. Orlov.

#### REFERENCES

- ALEXEJEVA, L. I. 1965. Stratigraphical review of the Proboscideans of Eopleistocene (according to the materials of the European south of the USSR). *V kn. "Chetvertichnyj period i ego istoria"*, "Nauka", 69-90.
- ALEXEJEVA, L. I. 1968. Die asiatischen Elemente in der Saugetierfauna des osteuropaischen Anthropogens. *Berichte Deutsch. Ges. Geol. Wiss., Geol. Palaeont.*, A. 13, 3, Berlin, 299-303.
- ALEXEJEVA, L. I. 1970. Early Quaternary Proboscideans of Europe. "Itogi nauki", *seria geologia*, "Nauka", 120-135.
- ALEXEJEVA, L. I., GARUTT, W. E. 1965. New data on the evolution of elephants of the genera *Archidiskodon*. *Bull. komis. po izuch. chetv. perioda AN SSSR*, N 30, M., "Nauka", 161-166.







- ARAMBOURG G. 1952. Note préliminaire sur quelques Elephantes fossiles de Berberie. *Bull. Mus. Nat. Hist. Nat.* Ser. 2, XXXIX, N2: 407-418.
- ARAMBOURG, G. 1970 a. Les Vertébrés du Pleistocène de l'Afrique du Nord. *Arch. Mus. Nat. Hist. Natur.* 7-c Ser., X, fasc. 1: 1-126.
- ARAMBOURG, G. 1970 b. Villafranchian correlations based on large Mammals. Proc. IV Session Commite Mediterr. Neogene Stratigraphy Bologna 1967, *Giornale di Geologia* (2), XXXV, fasc. 1.
- BAIGUSHEVA, T. S. 1964. The khaprovian fauna of Liventsovsky sand quarry (Rostov region). *Bull. kom. po izuch. chetv. perioda AN SSSR*, N 29: 44-50.
- BAIGUSHEVA, V. S. 1968. Paleontologic characteristics of mammalian fauna of the main upper pleistocene sections of the North-Eastern Priazovie (Liventzovka, Port-Katon). *Avto-ref. kand. diss., Rostov-na-Donu.* 3-18.
- BAIGUSHEVA, V. S. 1971. Fossil teriofauna of the Liventzovka sand-pit. V km. "Materialy po faunam antropogena SSSR", M., "Nauka". 5-29.
- BOGACHEV, V. V. 1924. New data on the Tertiary elephants history in the South-Eastern Russia. *I v. Azerb. Uniu-ta (za 1923-1924)*. N 3: 12-94.
- BURCHAK-ABRAMOVICH, N. I. and DJAFAROV, R.D. 1950. The data on the studies of Upper Tertiary Proboscidea of Trans-Caucasia. *Izv. Akad. nauk AzerbSSR*. N 2: 38-55.
- DUBROVO, I. A. 1964. Elephants of the genera *Archidiskodon* in the territory of the USSR. *Paleont. Journ. Akad. nauk SSSR*, N 3: 82-94.
- DUBROVO, I. A., BAIGUSHEVA, V. S. 1964. Specific composition of the elephants of Khaprovian faunistic complex (on the data of Liventsovsky quarry). *Bull. Mosk. ob-va ispyt. pripody*, tom XXXIX, otd. geol. N 5: 133-135.
- FRIANT, K. 1959. A propos du développement des molaires de l'*Elephas meridionalis* Nesti. *Ann. Soc. geol. Nord.* N2: 113-128.
- GABUNIA, L. K. 1961. On the mammalian remains from the middle Pliocene of the North Caucasus (Kosyakino). *Paleont. Journ. Akad. nauk SSSR*. N1: 163-135.
- GABUNIA, L. K., VEKUA, A. K. 1963. Fossil elephant from Taribana. *Izdvo Akad. nauk. Gruz. SSR, Tbilisi.* 3-68.
- GARUTT, V. E. 1954. Southern elephant—*Archidiskodon meridionalis* (Nesti) from the Pliocene of the northern coast of the Sea of Azov. *Tr. Kom. po izuch. chetv. perioda Akad. nauk SSSR*, X. V. 2: 1-77.
- GARUTT, V. E. 1964. Das Mammut, *Mammuthus primigenius* (Blum.). Die neue Bräun.—Bucherei, A, Ziemsen Verlag Wittenberg Lutherstadt.
- GARUTT, V. E. 1965. Fossil elephants of Siberia. *Tr. nauchno-issl. in-ta geologii Arktiki.* 143: 106-130.
- GARUTT, V. E., ALEXEJEVA, L. I. 1964. New data on the evolution of elephants genera *Archidiskodon*. *Tez. dokl. Vsesouzn. soveshch. po izuch. chetv. perioda, Novosibirsk.* 7-8.
- GROMOV, V. I. 1933. Problem of glation plurality in connection with the study of Quaternary mammals. *Probl. sov. geol.* 3 (7)
- GROMOV, V. I. 1948. Paleontological and archeological ground of the stratigraphy of the Quaternary continental deposits in the territory of the USSR (mammals, the Paleolith). *Tr. in-ta geol. nauk Acad. nauk SSSR*, vyp. 48, geol. serv. N17: 1-52.
- GROMOV, V. I. 1957. Stratigraphic scheme of the Quaternary deposits of the USSR and its correlation with foreign schemes. *Tez. dokl. Vsesojuzn. Mezhdved. soveshch. po izuch. chetv. perioda, Moskva.*
- GROMOV, V. I., MIRCHINK, G. F. 1936. Quaternary period and its fauna. *Zhivotnyi mir SSSR*. I: 639-640.
- KHOKHLOVKINA V. A. 1940. Terraces of the azovian coast between Rostov and Taganrog. *Tr. in-ta geol. nauk. Akad. nauk SSSR*, ser. geol., 8, vyp. 28.
- LEBEDEVA N. A. 1972. Anthropogene of Priazovie. *Tr. Geologicheskogo in-ta Akad. nauk SSSR, "Nauka"*, vyp. 125: 5-106.
- MACCAGNO, A. M. 1962. L'*Elephas meridionalis* Nesti di Contrada "Madonna della Strada", Scopito (L'Aquila). *Atti. Accad. Sci. Fis. Mat. Napoli, Mem.* Ser. 3, 4(1).
- NIKIFOROVA, K. V., ALEXEJEVA, L. I. 1959. On the boundary between the Tertiary and Quaternary systems according the data of mammalian fauna. *Tr. Geol. in-ta Akad. nauk SSSR*, vyp. 32: 7-21.
- NIKONOV, A. A., PENKOV, A. V. TROFIMOV, B. A. WANGENGEIM E. A. DMITRIEVA, E. L. SOTNIKOVA, M. V. 1971. New data on the geologia and mammalian fauna of the Upper Pliocene and Lower Pleistocene of Southern Tadzhikistan. *Bull. Mosk. obshch. ispyt. priir., otd. geol.*, N 6: 135.
- OSBORN, H. F. 1942. Proboscidea. *Amer. Mus. Nat. Hist.* II, 805-1675.
- RODZIANKO, G. N. 1967. Pliocene of the Lower Don and Lower Volga. V sb. "Materialy geol. issledovaniy na territorii Nyzhnego Dona" Nizhnej Volgi", Rostov-na-Donu. 172-283.
- RODZIANKO, G. N. 1970. The Neogene system. *Geologia SSSR, "Nedra", Moskva.* 46(1): 410-447.
- SCHAUB, S. 1948. Das gebiss der Elephanten. *Vorhandlungen der Naturforschenden Gesellschaft in Basel*, LIX.
- VERESCAGIN, N. K. 1959. The mammals of the Caucasus (the history of fauna formation). *Izd-vo Akad. nauk SSSR, M. -L.* -1-703.
- VISLOBOKOVA, I. A. 1974. On the new find of *Archidiskodon gromovi* Garutt et Alexejeva in the Lower Eopleistocene deposits of the South of West Siberia. *Bull. Kom. po izuch. chetv. prioda*, N 42: 171-176.
- WEITHOFER, K. 1890. Die fossilen Proboscidi der Arnoteles in Toscana. *Beitr. Pal. Osterr. -Ung.* Bd. VIII: 107-240.
- YANKOVA (BAIGUSHEVA), V. S. 1959. Paleontologic finds from Liventsovsky sand quarry. *Izv. Rost. obl. kraev. mus., Izd-vo Rost. un-ta*, N1(3): 41-49.

## EXPLANATION OF PLATES

## PLATE I

1. Skull of *A. gromovi* from Liventsovka, holotype. Coll. Rostov-on Don Museum of Regional Study, N 113; front view 1/5 natur. size.
2. Ditto, sid view, 1/5 natur. size.
3. Ditto, view from below. 1/2 nature. size.

## PLATE II

1. Last upper molar M<sup>3</sup> of *A. gromovi* from Liventsovka; view from above. Coll. GIN, N 270-10. 1/2 natur. size.
2. Ditto, side view, 1/2 natur. size.
3. First upper milk premolar dP<sup>2</sup> of *A. gromovi* from Khapry. Coll. GIN, N 300-125. Natur. size. View from above.
4. Ditto, side view, natur. size.